

Measurement of the Pion Form Factor in the Range 1.04 – 1.38 GeV with the CMD-2 Detector

V. M. Aulchenko^{a,b}, R. R. Akhmetshin^a, V. Sh. Banzarov^a, L. M. Barkov^{a,b}, N. S. Bashtovoy^a, D. V. Bondarev^{a,b},
 A. E. Bondar^a, A. V. Bragin^a, A. A. Valishev^a, N. I. Gabyshev^a, D. A. Gorbachev^a, A. A. Grebeniuk^a,
 D. N. Grigoriev^{a,b}, S. K. Dhawan^d, D. A. Epifanov^a, A. S. Zaitsev^{a,b}, S. G. Zverev^a, F. V. Ignatov^a, V. F. Kazanin^{a,b},
 S. V. Karpov^a, I. A. Koop^{a,b}, P. P. Krovovny^{a,b}, A. S. Kuzmin^{a,b}, I. B. Logashenko^{a,c}, P. A. Lukin^a, A. P. Lysenko^a,
 A. I. Milstein^{a,b}, K. Yu. Mikhailov^a, I. N. Nesterenko^{a,b}, M. A. Nikulin^{a,b}, A. V. Otboev^a, V. S. Okhapkin^a,
 E. A. Perevedentsev^{a,b}, A. A. Polunin^a, A. S. Popov^a, S. I. Redin^a, B. L. Roberts, N. I. Root^a, A. A. Ruban^a,
 N. M. Ryskulov^a, A. L. Sibidanov^a, V. A. Sidorov^a, A. N. Skrinsky^a, V. P. Smakhtin, I. G. Snopkov^a, E. P. Solodov^{a,b},
 J. A. Thompson^{e†}, G. V. Fedotov^{a,b}, B. I. Khazin^{a,b}, V. W. Hughes^{d†}, A. G. Shamov^a, Yu. M. Shatunov^a,
 B. A. Shwartz^{a,b}, S. I. Eidelman^{a,b}, Yu. V. Yudin^a

^a Budker Institute of Nuclear Physics, Russian Acad. Sci., Siberian Div., Novosibirsk, 630090, Russia

^b Novosibirsk State University, Novosibirsk, 630090, Russia

^c Boston University, Boston, MA 02215, USA

^d Yale University, New Haven, CT 06511, USA

^e University of Pittsburgh, Pittsburgh, PA 15260, USA

^f Weizmann Institute of Science, 76100, Rehovot, Israel

Published in JETP LETTERS Vol. 82 No. 12 (2005) 743-747

The cross section for the process $e^+e^- \rightarrow \pi^+\pi^-$ is measured in the c.m. energy range 1.04 ÷ 1.38 GeV from 995 000 selected collinear events including 860000 e^+e^- events, 82000 $\mu^+\mu^-$ events, and 33000 $\pi^+\pi^-$ events. The systematic and statistical errors of measuring the pion form factor are equal to 1.2 ÷ 4.2 and 5 ÷ 13 %, respectively.

PACS: 13.25.-k, 13.40.Gp, 13.66.-a, 29.30.-h

INTRODUCTION

A study of the cross section for the process $e^+e^- \rightarrow \pi^+\pi^-$ provides important information on the electromagnetic form factor of the pion, which describes its internal structure. Moreover, precision measurement of this cross section is necessary for calculating the anomalous magnetic moment of the muon $(g-2)_\mu$ [1] and its comparison with precision measurements, one of which was carried out recently at BNL [2]. Such comparison is an important test of the Standard Model.

EXPERIMENT

Measurements were performed at the VEPP-2M collider [3] with the CMD-2 general purpose detector (cryogenic magnetic detector), which combines the properties of a magnetic spectrometer and good calorimetry [4, 5]. The coordinates, emission angles, and momenta of charged particles are measured by the coordinate system of the detector, which consists of the drift and Z chambers located inside a thin (0.38 X_0) superconducting solenoid with a magnetic field of 1 T. Cylindrical and endcap electromagnetic calorimeters based on CsI and BGO scintillation crystals ensure measurement of

energy and photon emission angles and make it possible to separate electrons and hadrons. A range system is used to identify muons.

This work continues a cycle of precision measurements of hadron cross sections with the CMD-2 detector. The results of the pion form factor measurement in the energy range 0.61 ÷ 0.96 GeV were published in [6]. In this work, we present the results of the measurement of the form factor in the energy range 1.04 ÷ 1.38 GeV. A more detailed description of the data analysis was given in [7].

An integrated luminosity of 6 pb⁻¹ was collected in the experiment. For analysis, we selected 33 000 $\pi^+\pi^-$ events accumulated at 35 beam energy points from 520 to 690 MeV with a step of 5 MeV. The beam energy was controlled with an accuracy not worse than $\delta E/E \sim 10^{-3}$ using the magnetic field value in the VEPP-2M storage ring.

SELECTION OF COLLINEAR EVENTS

To separate the events $e^+e^- \rightarrow e^+e^-$, $e^+e^- \rightarrow \mu^+\mu^-$, and $e^+e^- \rightarrow \pi^+\pi^-$, the following conditions were used:

- One vertex with two tracks of particles with opposite charges was found in the drift chamber.
- The event vertex was located near the beam interaction point; i.e., $\rho_{vtx} = \min(\rho_{tr}^+, \rho_{tr}^-) < 0.15$ cm, where $\rho_{tr}^\pm$ is the minimum distance between the particle track and beam axis, and $|Z_{vtx}| < 10$ cm, where Z_{vtx} is the position of the vertex along the beam axis.
- Track collinearity conditions:
 $|\Delta\varphi| = |\pi - |\varphi^+ - \varphi^-|| < 0.15$, where $\varphi^\pm$ is the azimuth track angle;
 $|\Delta\theta| = |\pi - (\theta^+ + \theta^-)| < 0.25$, where $\theta^\pm$ is the polar track angle.
- Constraint on the solid angle of the event detection:
 $\theta_{min} < (\pi + \theta^- - \theta^+)/2 < \pi - \theta_{min}$, where $\theta_{min} = 1.1$.
- The mean momentum was bounded from above to reduce the cosmic particle background and from below to suppress $e^+e^- \rightarrow K^+K^-$ events:
 $E_{beam} + 150 \text{ MeV}/c > (p^+ + p^-)/2 >$
 $> \max(\sqrt{E_{beam}^2 - 494^2} \cdot 1.15 \text{ MeV}/c, 300 \text{ MeV}/c)$,
 where p^+ and p^- are the momenta of the positive and negative particles, respectively.

The main sources of the background for the process $e^+e^- \rightarrow \pi^+\pi^-$ are the $e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$, $e^+e^- \rightarrow \pi^+\pi^-\pi^0$, $e^+e^- \rightarrow K^+K^-$ reactions and cosmic particles. The physical background contribution to the pion form factor was calculated using experimental cross sections [8, 9] by taking into account the detection efficiency determined from the total simulation. The total contribution of these processes does not exceed 0.8% and is taken into account as a correction to the pion form factor according to Eq. (2). The number of cosmic particle background events was determined from the distribution of the event vertices over the distance from the beam interaction point.

EVENT SEPARATION

To determine the number of events of each process, we used two-dimensional distributions over the energy deposition in the CsI calorimeter (Fig. 1). The number of events (N_π , $N_\mu + N_e$) was determined by minimizing the maximum likelihood function:

$$\mathcal{L} = - \sum_{events} \ln \left(\sum_i N_i \cdot f_i(E^+, E^-) \right) + \sum_i N_i,$$

where f_i is the probability density function for events of a given type ($\pi, \mu, e, cosmic$). Electrons and positrons initiate an electromagnetic shower in a calorimeter and thereby noticeably differ from other charged particles




Fig. 1. Distribution of collinear particles over the energy deposition in the calorimeter.

in their energy deposition. The energy depositions of muons, cosmic particles, and pions with only ionization losses are almost identical. For this reason, to determine the number of muons, we used additional information on the ratio of the cross section for muon production to the cross section for $e^+e^- \rightarrow e^+e^-$ as obtained from the QED calculation with the inclusion of the resolutions and detection efficiencies in the detector.

DETERMINATION OF THE PION FORM FACTOR

The cross section for the $e^+e^- \rightarrow \pi^+\pi^-$ process integrated over the detection solid angle is given by the expression:

$$\begin{aligned} \sigma_{\pi^+\pi^-} &= \sigma_{\pi^+\pi^-}^0 \cdot |F_\pi|^2 \\ &= \frac{\pi\alpha^2}{3s} \left(1 - \frac{4m_\pi^2}{s}\right)^{\frac{3}{2}} \frac{3 \cos \theta_{min} - \cos^3 \theta_{min}}{2} \cdot |F_\pi|^2, \end{aligned} \quad (1)$$

where $\sigma_{\pi^+\pi^-}^0$ is the cross section calculated under the assumption of the absence of the internal structure of the pion. The contribution of vacuum polarization to the photon propagator is also included in the form factor.

At each energy point, the form factor is calculated as:

$$|F_\pi|^2 = \frac{N_{\pi^+\pi^-}}{N_{e^+e^-} + N_{\mu^+\mu^-}} \times \frac{\sigma_{e^+e^-}^0 \cdot (1 + \delta_{e^+e^-}) \cdot \varepsilon_{e^+e^-} + \sigma_{\mu^+\mu^-}^0 \cdot (1 + \delta_{\mu^+\mu^-}) \cdot \varepsilon_{\mu^+\mu^-}}{\sigma_{\pi^+\pi^-}^0 \cdot (1 + \delta_{\pi^+\pi^-}) \cdot \varepsilon_{\pi^+\pi^-} \cdot (1 - \Delta_{\pi \text{ loss}}) - \Delta_{3\pi, 4\pi, K^+K^-}}, \quad (2)$$

Here, $N_{\pi^+\pi^-}/(N_{e^+e^-} + N_{\mu^+\mu^-})$ is the ratio of the number of detected pions to the number of muons and electrons as obtained in the minimization procedure, σ_{ii}^0 is the Born cross section in the lowest order of perturbation theory, δ_{ii} is the radiative correction, ε_{ii} is the detection efficiency, $\Delta_{3\pi, 4\pi, K^+K^-}$ is the correction for background processes, and Δ_{loss} is the correction for the pion loss at the vacuum chamber wall and drift chamber material due to nuclear interactions. The Δ_{loss} correction was determined from simulation by comparing the number of selected pions with the inclusion and exclusion of nuclear interactions. The correction value was equal to $0.8 \div 1.2\%$.

Collinear events were selected using only information from the drift chamber. For this reason, by selecting desired (test) events from the CsI calorimeter and checking whether the reconstructed tracks are in the drift chamber, one can determine the event reconstruction efficiency. The reconstruction efficiency was $97 \div 98\%$. According to Eq. (2), only the difference between the detection efficiencies for different processes, which was $0.16 \pm 0.09\%$ between electrons and muons, is important for the determination of the pion form factor.

The radiative corrections for the processes $e^+e^- \rightarrow e^+e^-$, $e^+e^- \rightarrow \mu^+\mu^-$, and $e^+e^- \rightarrow \pi^+\pi^-$ are calculated using a procedure that was developed in [10] and was based on the formulas from [11, 12]. The probability of the emission of many photons along the initial and final particles, the emission of one photon at a large angle, and the vacuum polarization contribution to the photon propagator are taken into account in these formulas. Since the vacuum polarization is usually included in the definition of the form factor, its contribution was not included in the radiative correction for the process $e^+e^- \rightarrow \pi^+\pi^-$. According to [10], the accuracy of the calculation of the radiative corrections is estimated as 0.2% for all the processes. The number of selected collinear events depends on the angular and momentum resolutions of the drift chamber. In order to include them in the calculation for radiative corrections, the emission angles and momenta of particles were additionally simulated according to the experimental resolution and then the selection criteria were imposed. For

Table 1. Experimental pion form factor $|F_\pi|^2$. Only the statistical error is given

E , MeV	$ F_\pi ^2$	E , MeV	$ F_\pi ^2$
490.0	3.596 ± 0.163	605.7	1.069 ± 0.082
520.0	2.598 ± 0.134	610.0	0.989 ± 0.075
525.0	2.262 ± 0.112	615.0	1.069 ± 0.088
530.0	2.185 ± 0.135	620.0	0.988 ± 0.081
535.0	2.295 ± 0.130	625.0	0.794 ± 0.064
540.0	1.884 ± 0.119	630.0	0.696 ± 0.063
545.0	2.120 ± 0.110	635.0	0.719 ± 0.057
550.0	1.704 ± 0.120	640.0	0.693 ± 0.052
555.0	1.641 ± 0.106	645.0	0.571 ± 0.042
560.0	1.449 ± 0.146	650.0	0.640 ± 0.046
565.0	1.683 ± 0.103	655.0	0.570 ± 0.050
570.0	1.531 ± 0.088	660.0	0.483 ± 0.054
575.0	1.374 ± 0.150	665.0	0.460 ± 0.040
580.0	1.386 ± 0.087	670.0	0.524 ± 0.062
585.0	1.197 ± 0.115	675.0	0.347 ± 0.049
590.0	1.200 ± 0.088	680.0	0.357 ± 0.040
595.0	1.014 ± 0.093	685.0	0.424 ± 0.078
600.0	0.983 ± 0.079	690.0	0.338 ± 0.032

the $e^+e^- \rightarrow e^+e^-$ process, the bremsstrahlung energy losses of electrons and positrons on the vacuum chamber wall and first 10 cm of the drift chamber were taken into account.

Table 1 presents the form factor at each energy point.

SYSTEMATIC ERROR

The main contributions to the systematic error are listed in Table 2. The systematic error increases with energy, because the error in the number of muons directly contributes to the error in the number of pions, and the ratio of the number of muons to the number of pions increases from 1 to 7 when the c.m. energy increases from 1 to 1.38 GeV. The total systematic error is equal to $1.2 \div 4.2\%$ and does not exceed one third of the statistical error at each experimental point.

One of the tests of the separation procedure was performed using the simulation of e^+e^- , $\mu^+\mu^-$, and $\pi^+\pi^-$ events. The simulation data were analyzed with the inclusion of the following corrections: pion loss due to the nuclear interaction, the energy losses of electrons at the

Table 2. Various contributions to the systematic error in $|F_\pi|^2$. The given range corresponds to the scanned energy range

Error source	Error $\sqrt{s} = 1.04 \div 1.38$ GeV
Detection solid angle	$0.2 \div 0.5$ %
Detection efficiency	$0.5 \div 2$ %
Pion loss	0.2 %
Bremsstrahlung e^+e^-	$0.05 \div 1.7$ %
Radiative corrections	$0.5 \div 2$ %
Background events	$0.6 \div 1.6$ %
Energy calibration	$0.7 \div 1.1$ %
Particle separation procedure	$0.2 \div 1.5$ %
	$1.2 \div 4.2$ %
Statistical error at the point	$5 \div 13$ %

vacuum-chamber wall, and the resolution of the drift chamber while calculating radiative corrections. The calculated difference between the detection efficiencies for e^+e^- and $\mu^+\mu^-$, $\pi^+\pi^-$ in the simulation was equal to $\varepsilon_{MIP} - \varepsilon_{e^+e^-} = 0.189 \pm 0.004$ % in good agreement with the measured value. The difference between the form factor obtained and the form factor used in the simulation varies from 0.2 to 1.5 % in dependence on the energy. The difference at the highest energy was equal to 1.5 % and consisted of 1 % for the separation procedure and 0.5 % characterizing the systematic error in the inclusion of the above corrections.

DISCUSSION

Figure 2 shows the results, which are in good agreement with the data obtained in the previous experiments with the detectors OLYA[13], DM1[14], DM2[15], BCF[16], ACO[17]. The form factor in this energy range was measured in detail only in the experiment with the OLYA detector with a systematic error of $10 \div 15$ %. The experimental energy dependence of the form factor is well reproduced in the framework of the vector-meson dominance model by the sum of the amplitudes of the $\rho(770)$, $\rho(1450)$, $\rho(1700)$, ω and ϕ mesons [6]:

$$|F_\pi(s)|^2 = \left| \left(\text{BW}_{\rho(770)}^{\text{GS}}(s) \cdot \left(1 + \delta_\omega \frac{s}{m_\omega^2} \text{BW}_\omega(s) + \delta_\phi \frac{s}{m_\phi^2} \text{BW}_\phi(s) \right) + \beta \text{BW}_{\rho(1450)}^{\text{GS}}(s) + \gamma \text{BW}_{\rho(1700)}^{\text{GS}}(s) \right) / (1 + \beta + \gamma) \right|^2, \quad (3)$$




Fig. 2. Experimental data obtained for the pion form factor $|F_\pi|^2$ in this work in comparison with other experiments.

Here, $\text{BW}_\rho^{\text{GS}}(s)$ is the meson parametrization in the Gounaris-Sakurai model [18]; $\text{BW}_\omega(s)$ and $\text{BW}_\phi(s)$ are the parametrization of the ω and ϕ resonances, respectively, which were represented by the relativistic Breit-Wigner form due to a small width; $\delta_\omega, \delta_\phi$, β and γ are the model parameters describing the relative contributions of the $\rho - \omega$ and $\rho - \phi$ interferences and $\rho(1450)$ and $\rho(1700)$ states, respectively. In order to determine the model parameters, it is necessary to use all the available data on the form factor in the energy range $\sqrt{s} = 0.36 \div 3.7$ GeV, which will be done in a future work with analysis of all the information accumulated at the CMD-2 detector in the energy range from 0.37 to 1.38 GeV.

CONCLUSIONS

In this work, the cross section for the process $e^+e^- \rightarrow \pi^+\pi^-$ was measured in the c.m. energy range $1.04 \div 1.38$ GeV with the world best accuracy. The systematic and statistical errors of the measurement are equal to $1.2 \div 4.2$ % and $5 \div 13$ %, respectively. The measured cross section agrees well with the results of the previous experiments.

This work was supported by the Russian Foundation for Basic Research (project nos. 03-02-16477, 03-0216280, 03-02-16843, 04-02-16217, 04-02-16223, and 04-02-16434).

1. M. Davier and W. J. Marciano, Ann. Rev. Nucl. Part. Sci. **54**, 115 (2004).
2. G. W. Bennett et al., Phys. Rev. Lett. **92**, 161802 (2004).

3. *V. V. Anashin, I. B. Vasserman, V. G. Veshcherevich et al.*, Preprint Budker INP **84-114**, Novosibirsk, 1984.
4. *G. A. Aksenov, V. M. Aulchenko, L. M. Barkov et al.*, Preprint Budker INP **85-118**, Novosibirsk, 1985.
5. *E. V. Anashkin, V. M. Aulchenko, S. E. Baru et al.*, ICFA Instrumentation Bulletin **5**, 18 (1988).
6. *R. R. Akhmetshin et al.*, Phys. Lett. B **527**, 161 (2002),
R. R. Akhmetshin et al., Phys. Lett. B **578**, 285 (2004).
7. *V. M. Aulchenko, R. R. Akhmetshin, V. Sh. Banzarov et al.*, Preprint Budker INP **2005-29**, Novosibirsk, 2005.
8. *R. R. Akhmetshin et al.*, Phys. Lett. B **489**, 125 (2000).
9. *R. R. Akhmetshin et al.*, Phys. Lett. B **466**, 392 (1999).
10. *A. B. Arbuzov, G. V. Fedotovitch, F. V. Ignatov et al.*, Preprint Budker INP **04-70**, Novosibirsk, 2004.
11. *A. B. Arbuzov, E. A. Kuraev, V. A. Astakhov et al.*, JHEP **10**, 006 (1997).
12. *A. B. Arbuzov, E. A. Kuraev, G. V. Fedotovitch et al.*, JHEP **10**, 001 (1997).
13. *L. M. Barkov et al.*, Nucl. Phys. B **256**, 365 (1985).
14. *A. Quenzer et al.*, Phys. Lett. B **76**, 512 (1978).
15. *D. Bisello et al.*, Phys. Lett. B **220**, 321 (1989).
16. *D. Bollini et al.*, Lett. Nuovo Cim. **14**, 418 (1975).
17. *G. Cosme et al.*, Preprint LAL-1287, Orsay, 1976.
18. *G. J. Gounaris and J. J. Sakurai*, Phys. Rev. Lett. **21**, 244 (1968).